

Philippines - The probabilities of a hip fracture in men

The following tables give the 10-year probability (%) of a hip fracture according to body mass index (BMI kg/m²), the number of clinical risk factors (CRF) and age. Each table provides a mean estimate and a range, based on the epidemiology of Philippines. The range is not a confidence interval, but because the weight of different risk factors varies, is a true range.

Age = 50 years

Number of CRFs	BMI (kg/m ²)						
	15	20	25	30	35	40	45
0	0.2	0.1	0.1	0.1	0.0	0.0	0.0
1	0.4 (0.3-0.8)	0.2 (0.2-0.5)	0.1 (0.1-0.3)	0.1 (0.1-0.2)	0.1 (0.1-0.2)	0.1 (0.0-0.1)	0.0 (0.0-0.1)
2	0.8 (0.4-1.6)	0.5 (0.2-1.0)	0.3 (0.1-0.6)	0.2 (0.1-0.4)	0.2 (0.1-0.3)	0.1 (0.1-0.3)	0.1 (0.1-0.2)
3	1.4 (0.6-2.8)	0.9 (0.3-1.7)	0.5 (0.2-1.0)	0.4 (0.2-0.8)	0.3 (0.1-0.6)	0.2 (0.1-0.5)	0.2 (0.1-0.4)
4	2.5 (1.0-4.2)	1.5 (0.6-2.6)	0.9 (0.4-1.5)	0.7 (0.3-1.2)	0.5 (0.2-0.9)	0.4 (0.2-0.7)	0.3 (0.1-0.5)
5	4.4 (1.9-5.7)	2.7 (1.2-3.6)	1.6 (0.7-2.1)	1.2 (0.6-1.7)	1.0 (0.4-1.3)	0.7 (0.3-1.0)	0.6 (0.3-0.8)
6	7.3	4.7	2.8	2.2	1.7	1.3	1.0

Ten year probability of hip fractures (%) according to BMI, the number of clinical risk factors (CRF) and age in men from Phillipines.

Age = 55 years

Number of CRFs	BMI (kg/m ²)						
	15	20	25	30	35	40	45
0	0.3	0.2	0.1	0.1	0.1	0.1	0.0
1	0.6 (0.4-1.1)	0.4 (0.3-0.7)	0.2 (0.2-0.4)	0.2 (0.1-0.3)	0.1 (0.1-0.2)	0.1 (0.1-0.2)	0.1 (0.1-0.1)
2	1.1 (0.6-2.0)	0.7 (0.4-1.3)	0.4 (0.2-0.8)	0.3 (0.2-0.6)	0.2 (0.1-0.5)	0.2 (0.1-0.4)	0.1 (0.1-0.3)
3	1.9 (0.9-3.5)	1.2 (0.5-2.2)	0.7 (0.3-1.4)	0.6 (0.3-1.0)	0.4 (0.2-0.8)	0.3 (0.2-0.6)	0.2 (0.1-0.5)
4	3.2 (1.5-5.2)	2.1 (1.0-3.4)	1.3 (0.6-2.1)	1.0 (0.4-1.6)	0.8 (0.3-1.2)	0.6 (0.3-0.9)	0.4 (0.2-0.7)
5	5.4 (2.7-6.8)	3.6 (1.8-4.6)	2.2 (1.1-2.9)	1.7 (0.9-2.2)	1.3 (0.7-1.7)	1.0 (0.5-1.3)	0.8 (0.4-1.0)
6	8.8	6.0	3.8	2.9	2.2	1.7	1.3

Age = 60 years

Number of CRFs	BMI (kg/m ²)						
	15	20	25	30	35	40	45
0	0.5	0.3	0.2	0.1	0.1	0.1	0.1
1	0.8 (0.6-1.4)	0.5 (0.4-0.9)	0.3 (0.2-0.5)	0.3 (0.2-0.4)	0.2 (0.1-0.3)	0.2 (0.1-0.3)	0.1 (0.1-0.2)
2	1.4 (0.8-2.4)	1.0 (0.5-1.6)	0.6 (0.3-1.1)	0.5 (0.3-0.8)	0.4 (0.2-0.6)	0.3 (0.2-0.5)	0.2 (0.1-0.4)
3	2.4 (1.2-4.1)	1.6 (0.8-2.9)	1.0 (0.5-1.8)	0.8 (0.4-1.4)	0.6 (0.3-1.1)	0.5 (0.2-0.8)	0.4 (0.2-0.6)
4	3.9 (2.0-6.2)	2.7 (1.5-4.3)	1.8 (0.9-2.8)	1.4 (0.7-2.2)	1.0 (0.5-1.7)	0.8 (0.4-1.3)	0.6 (0.3-1.0)
5	6.3 (3.6-8.0)	4.5 (2.6-5.6)	3.0 (1.7-3.8)	2.3 (1.3-2.9)	1.8 (1.0-2.2)	1.3 (0.8-1.7)	1.0 (0.6-1.3)
6	9.8	7.3	5.0	3.8	2.9	2.2	1.7

Ten year probability of hip fractures (%) according to BMI, the number of clinical risk factors (CRF) and age in men from Phillipines.

Age = 65 years

Number of CRFs	BMI (kg/m ²)						
	15	20	25	30	35	40	45
0	0.6	0.4	0.3	0.2	0.2	0.1	0.1
1	1.0 (0.8-1.6)	0.7 (0.6-1.1)	0.5 (0.4-0.7)	0.4 (0.3-0.5)	0.3 (0.2-0.4)	0.2 (0.2-0.3)	0.2 (0.1-0.2)
2	1.7 (1.0-2.7)	1.2 (0.8-1.9)	0.8 (0.5-1.3)	0.6 (0.4-1.0)	0.5 (0.3-0.7)	0.4 (0.2-0.6)	0.3 (0.2-0.4)
3	2.7 (1.6-4.4)	2.0 (1.1-3.3)	1.4 (0.8-2.3)	1.0 (0.6-1.7)	0.8 (0.4-1.3)	0.6 (0.3-1.0)	0.5 (0.3-0.8)
4	4.3 (2.5-6.6)	3.2 (1.9-4.9)	2.2 (1.4-3.4)	1.7 (1.0-2.6)	1.3 (0.8-2.0)	1.0 (0.6-1.5)	0.7 (0.4-1.2)
5	6.7 (4.3-8.5)	5.1 (3.3-6.4)	3.6 (2.4-4.5)	2.8 (1.8-3.4)	2.1 (1.4-2.6)	1.6 (1.0-2.0)	1.2 (0.8-1.5)
6	10	7.9	5.8	4.4	3.4	2.6	1.9

Age = 70 years

Number of CRFs	BMI (kg/m ²)						
	15	20	25	30	35	40	45
0	0.8	0.6	0.4	0.3	0.2	0.2	0.1
1	1.3 (0.9-1.6)	1.0 (0.7-1.2)	0.7 (0.5-0.9)	0.5 (0.4-0.7)	0.4 (0.3-0.5)	0.3 (0.2-0.4)	0.2 (0.2-0.3)
2	2.2 (1.3-3.4)	1.7 (1.1-2.6)	1.2 (0.8-1.8)	0.9 (0.6-1.4)	0.7 (0.5-1.0)	0.5 (0.3-0.8)	0.4 (0.3-0.6)
3	3.6 (2.0-5.8)	2.8 (1.7-4.5)	2.1 (1.3-3.2)	1.6 (1.0-2.4)	1.2 (0.7-1.8)	0.9 (0.5-1.4)	0.7 (0.4-1.1)
4	5.9 (3.6-8.6)	4.7 (2.9-7.0)	3.6 (2.3-5.3)	2.7 (1.7-4.1)	2.0 (1.3-3.1)	1.5 (1.0-2.3)	1.2 (0.7-1.8)
5	9.3 (7.3-13)	7.6 (6.0-10)	6.0 (4.7-8.0)	4.5 (3.6-6.1)	3.4 (2.7-4.6)	2.6 (2.0-3.5)	2.0 (1.5-2.7)
6	14	12	9.7	7.4	5.6	4.3	3.2

Ten year probability of hip fractures (%) according to BMI, the number of clinical risk factors (CRF) and age in men from Phillipines.

Age = 75 years

Number of CRFs	BMI (kg/m ²)						
	15	20	25	30	35	40	45
0	1.0	0.8	0.6	0.4	0.3	0.3	0.2
1	1.8 (1.1-3.3)	1.4 (0.9-2.7)	1.1 (0.7-2.0)	0.8 (0.5-1.5)	0.6 (0.4-1.2)	0.5 (0.3-0.9)	0.4 (0.2-0.7)
2	3.2 (1.6-5.8)	2.6 (1.3-4.7)	2.0 (1.1-3.7)	1.5 (0.8-2.8)	1.2 (0.6-2.1)	0.9 (0.5-1.6)	0.6 (0.3-1.2)
3	5.4 (2.4-9.8)	4.5 (2.0-8.1)	3.6 (1.7-6.3)	2.7 (1.3-4.8)	2.1 (0.9-3.7)	1.5 (0.7-2.8)	1.2 (0.5-2.1)
4	8.9 (4.2-14)	7.6 (3.6-12)	6.2 (2.9-9.7)	4.7 (2.2-7.4)	3.6 (1.7-5.6)	2.7 (1.2-4.3)	2.0 (0.9-3.2)
5	14 (7.4-19)	12 (6.3-17)	10 (5.2-14)	7.8 (4.0-11)	6.0 (3.0-8.4)	4.5 (2.2-6.4)	3.4 (1.7-4.8)
6	22	19	16	13	9.7	7.4	5.6

Age = 80 years

Number of CRFs	BMI (kg/m ²)						
	15	20	25	30	35	40	45
0	1.3	1.0	0.8	0.6	0.5	0.4	0.3
1	2.3 (1.4-4.3)	1.9 (1.2-3.6)	1.5 (1.0-2.9)	1.1 (0.7-2.2)	0.9 (0.6-1.6)	0.6 (0.4-1.2)	0.5 (0.3-0.9)
2	3.9 (2.1-7.3)	3.3 (1.8-6.2)	2.7 (1.5-5.0)	2.0 (1.1-3.8)	1.5 (0.8-2.8)	1.1 (0.6-2.1)	0.8 (0.5-1.6)
3	6.5 (3.1-11)	5.5 (2.7-9.3)	4.6 (2.2-7.6)	3.4 (1.7-5.8)	2.6 (1.2-4.4)	1.9 (0.9-3.3)	1.5 (0.7-2.5)
4	10 (4.8-16)	8.8 (4.1-13)	7.5 (3.5-11)	5.7 (2.6-8.6)	4.3 (1.9-6.5)	3.2 (1.4-4.9)	2.4 (1.1-3.7)
5	16 (8.2-21)	14 (7.0-18)	12 (5.9-16)	9.1 (4.5-12)	6.9 (3.4-9.5)	5.2 (2.5-7.3)	4.0 (1.9-5.5)
6	23	21	18	14	11	8.3	6.3

Ten year probability of hip fractures (%) according to BMI, the number of clinical risk factors (CRF) and age in men from Phillipines.

Age = 85 years

Number of CRFs	BMI (kg/m ²)						
	15	20	25	30	35	40	45
0	1.6	1.3	1.0	0.8	0.6	0.4	0.3
1	2.9 (1.9-5.4)	2.4 (1.5-4.5)	1.9 (1.2-3.5)	1.4 (0.9-2.7)	1.1 (0.7-2.0)	0.8 (0.5-1.5)	0.6 (0.4-1.1)
2	5.0 (2.7-9.2)	4.1 (2.3-7.6)	3.3 (1.9-6.1)	2.5 (1.4-4.6)	1.9 (1.0-3.5)	1.4 (0.8-2.6)	1.1 (0.6-2.0)
3	8.2 (4.1-14)	6.9 (3.4-11)	5.6 (2.8-9.2)	4.3 (2.1-7.1)	3.2 (1.6-5.4)	2.4 (1.2-4.1)	1.8 (0.9-3.1)
4	13 (6.2-19)	11 (5.3-16)	9.2 (4.4-14)	7.1 (3.3-10)	5.4 (2.5-8.0)	4.1 (1.8-6.1)	3.1 (1.4-4.6)
5	20 (10-26)	17 (8.9-23)	15 (7.5-19)	11 (5.7-15)	8.6 (4.3-12)	6.6 (3.2-9.0)	5.0 (2.4-6.9)
6	28	25	22	17	13	10	7.9

Age = 90 years

Number of CRFs	BMI (kg/m ²)						
	15	20	25	30	35	40	45
0	1.5	1.2	0.9	0.7	0.5	0.4	0.3
1	2.7 (1.8-5.1)	2.2 (1.4-4.1)	1.7 (1.1-3.2)	1.3 (0.9-2.4)	1.0 (0.6-1.8)	0.7 (0.5-1.4)	0.5 (0.4-1.0)
2	4.7 (2.6-8.7)	3.8 (2.1-7.1)	3.0 (1.7-5.6)	2.3 (1.3-4.2)	1.7 (1.0-3.2)	1.3 (0.7-2.4)	1.0 (0.5-1.8)
3	7.8 (3.9-13)	6.5 (3.2-11)	5.2 (2.6-8.5)	3.9 (2.0-6.5)	3.0 (1.5-4.9)	2.2 (1.1-3.7)	1.7 (0.8-2.8)
4	12 (6.0-18)	10 (5.0-15)	8.6 (4.0-13)	6.5 (3.0-9.7)	5.0 (2.3-7.4)	3.8 (1.7-5.6)	2.8 (1.3-4.3)
5	19 (10-25)	16 (8.4-21)	14 (6.9-18)	11 (5.3-14)	8.1 (4.0-11)	6.1 (3.0-8.4)	4.6 (2.2-6.4)
6	27	24	21	16	13	9.8	7.5