

NMR STUDIES OF STRUCTURE AND FUNCTION OF BIOLOGICAL MACROMOLECULES

Nobel Lecture, December 8, 2002

by

KURT WÜTHRICH

Eidgenössische Technische Hochschule Zürich, CH-8093 Zürich, Switzerland, and The Scripps Research Institute, 10550 N. Torrey Pines Rd., La Jolla, CA 92037, USA.

INTRODUCTION

Nuclear magnetic resonance (NMR) spectroscopy is unique among the methods available for three-dimensional structure determination of proteins and nucleic acids at atomic resolution, since the NMR data can be recorded in solution. Considering that body fluids such as blood, stomach liquid and saliva are protein solutions where these molecules perform their physiological functions, knowledge of the molecular structures in solution is highly relevant. In the NMR experiments, solution conditions such as the temperature, pH and salt concentration can be adjusted so as to closely mimic a given physiological fluid. Conversely, the solutions may also be changed to quite extreme non-physiological conditions, for example, for studies of protein denaturation. Furthermore, in addition to protein structure determination, NMR applications include investigations of dynamic features of the molecular structures, as well as studies of structural, thermodynamic and kinetic aspects of interactions between proteins and other solution components, which may either be




Figure 1. NMR structure of the *Antennapedia* homeodomain (1). A bundle of 20 superimposed conformers represents the polypeptide backbone. For the polypeptide segment 7–59 the tight fit of the bundle indicates that the structure is defined with high precision, whereas the two chain ends are disordered.




Figure 2. NMR structure of the complex formed between the *Antennapedia* homeodomain (blue, with functionally important residues in red, yellow, green and brown) and its operator DNA (yellow and red) (2).

other macromolecules or low molecular weight ligands. Again, for these supplementary data it is of keen interest that they can be measured directly in solution.

The NMR structure of the *Antennapedia* homeodomain (Figure 1) illustrates one of the exciting features of being able to perform structural studies in solution. The polypeptide chain in this protein is only partially folded, with both chain ends showing pronounced disorder (1). In the complex with its operator DNA (Figure 2) the N-terminal chain end is located in the minor groove of the DNA, where it adopts a well-defined structure (2). This mode of intermolecular recognition illustrates the functional importance of partially structured polypeptide chains. Mammalian prion proteins are an even more striking example of partial polypeptide folding, since the three-dimensional structure of the benign “cellular” form (PrP^C) includes a flexibly disordered 100-residue tail linked to the N-terminal end of a globular domain (Figures 3 and 4) (3). Considering that the mechanism of transformation of PrP^C into the aggregated, disease-related form of mammalian prion proteins is still subject to speculation, the observation of this flexible tail has been highly intriguing.

Partially folded polypeptide chains are usually difficult to crystallize. Furthermore, if crystals can be obtained, the chain segments that are disordered in solution will either be ordered by intermolecular contacts in the crystal lattice, or they will not be visible by diffraction methods. As a consequence, NMR has in many cases been the only method capable of providing structural information on partially folded polypeptides. Although a standard protocol for NMR structure determination provides a static picture of the unstructured chain segments (Figures 1 and 4) (4), additional NMR experi-




Figure 3. NMR structure of the bovine prion protein (3). In the C-terminal globular domain of residues 126–230, α -helices are green, an antiparallel β -sheet is blue, and non-regular secondary structure is yellow; the “unstructured” N-terminal tail of residues 23–125 is white.

ments can provide information on the frequencies of the rate processes that mediate transitions between discrete states of the molecule within the conformation space spanned by the static bundle of NMR-conformers (5, 6).

The ability of NMR techniques to characterize macromolecular structures and their intermolecular interactions with high spatial and temporal resolution has long attracted keen interest. This article reports on experience gained with NMR studies of proteins in my laboratory during the past 35 years (7).




Figure 4. Visual impression of the variation of the bovine prion protein structure (Figure 3) during a time period of about 1 nanosecond. The superposition of 20 snapshots illustrates that the globular domain maintains its mean geometry, whereas the tail undergoes large-scale changes with time.

NMR SPECTRA OF PROTEINS IN SOLUTION

When I joined the field of biomacromolecular NMR spectroscopy in 1967, important qualitative NMR features of amino acids and proteins had already been noted and tentatively rationalized (8–12). Most important, it had been well documented that the spectrum of a globular protein is more complex than the sum of the NMR lines from the constituent amino acid residues in the polypeptide chain (Figure 5), and the differences between the ^1H NMR spectra of folded and unfolded (“random coil”) forms of a polypeptide chain had been tentatively explained by different interactions with the solvent (Figure 6). The spectral analysis was primarily focused on the positions of the individual NMR lines in the ^1H NMR spectrum, as given by the “chemical shift”, δ , in parts per million (ppm) relative to a reference compound (Figure 5). Although the chemical shift is primarily determined by the covalent structure of the amino acid residue, it can also be significantly affected by the interactions with the solvent. Therefore, the exclusion of the solvent water from the interior of a globular protein (Figure 6) causes the chemical shifts of the core residues to be different from those of the water-exposed amino acid residues, so that even NMR lines originating from multiple residues of the same amino acid type can be distinguished. This “conformation-dependent chemical shift dispersion” was found to be sufficiently large to enable ^1H NMR studies of protein denaturation (Figure 7) (12). This then indicated the exciting prospect of using NMR for detailed studies of protein folding, and in particular for distinguishing between two-state and multiple-state folding and unfolding transitions. For me, the observations and ideas illustrated by Figures 5 to 7 also showed that one would need the ability of resolving and analyzing highly complex NMR spectra (upper trace in Figure 5) in order to obtain information about physiologically active, folded forms of proteins. This was a clearly defined problem that appeared sufficiently challenging to be attractive for my own research.




Figure 5. One-dimensional (1D) ^1H NMR spectra of the small protein bovine pancreatic trypsin inhibitor (BPTI, $M \approx 6000$). Top: Experimental spectrum of folded, active BPTI in a freshly prepared $^2\text{H}_2\text{O}$ -solution. Bottom: Simulated spectrum for the unfolded, random coil form of the BPTI polypeptide chain.




Figure 6. Scheme illustrating that the solvent (blue) has free access to all parts of a random coil polypeptide chain (rc), whereas it is excluded from the core of a folded globular protein.

In attempts at rationalizing in more detail the observations of Figures 5 and 7, it had been suggested that local magnetic fields generated by “ring currents” in the aromatic amino acid side chains (Figure 8) should cause outstandingly large conformation-dependent chemical shifts of hydrogen atoms located near the rings in the three-dimensional protein structure. This was qualitatively confirmed by comparison of the largest observed deviations from random coil chemical shifts (4) in hen egg white lysozyme (Figure 7) with calculations of the “ring current shifts” based on the then-available low-resolution crystal structure of this protein (12). It had also been pointed out that hemoproteins could be expected to have particularly large ring current shifts for hydrogen atoms located near the heme groups (Figure 9) (10,12).

For my start in the field of NMR with proteins, the prediction of unique, large heme ring current shifts seemed to be an attractive feature of hemo-




Figure 7. Low-field and high-field regions of 1D ^1H NMR spectra of the protein lysozyme at different temperatures, reflecting the transition from the folded form at 56 °C to the random coil form of the polypeptide chain at 73 °C (220 MHz, H_2O -solution) (adapted from ref. 12).




Figure 8. Local "ring current field" around aromatic rings in solution induced by an external, static magnetic field. The shape of the ring current field is indicated by the red double-cone and by broken magnetic-field-lines. The minus-sign indicates that the NMR lines of hydrogen atoms located inside the cone in the three-dimensional protein structure are shifted "upfield" (to the right in the spectra of Figure 5), whereas for atoms outside of the cone the shifts are "downfield".

proteins, and the presence of the heme iron (Figure 9) was appealing in view of my background in inorganic chemistry. As an additional advantage, low-resolution crystal structures for several hemoproteins were already available in 1967. It turned out that for several years nearly all my research projects were focused on the heme iron and its coordinatively linked ligands, and on the non-bonding interactions of the heme groups with their immediate environment in the hemoproteins.




Figure 9. Chemical structure of heme c, which is the prosthetic group in cytochromes c. The red arrows connect groups of hydrogen atoms in the covalent structure of heme c that are separated by a sufficiently short distance to be connected by nuclear Overhauser effects (NOE).




Figure 10. 1D ^1H NMR spectrum (360 MHz, 52°C , solvent $^2\text{H}_2\text{O}$) of horse ferrocyanochrome c ($M \approx 12000$).

As anticipated, the large heme ring current field generates a small number of well-separated resonance lines in the spectra of folded, globular hemoproteins (Figure 10). For the paramagnetic states of hemoproteins, additional well-resolved lines result from interactions with the unpaired electrons of the heme iron. Although they represented less than 3% of all the hydrogen atoms in any given hemoprotein, the well-resolved resonances were of crucial importance, because these lines could be selectively irradiated in 1D ^1H NMR experiments. For example, performing some simple ligand exchange and redox reactions with cytochrome c (Figure 11) resulted in the identification of one of the axial ligands bound to the heme iron, which had not been seen in the low resolution X-ray crystal structure available in 1969.

The experiments in Figure 11 (13) started with the oxidised form of cytochrome c. As was then learned through this study (see below), ferricyto-




Figure 11. Three chemical reaction steps with cytochrome c that were used to identify methionine as one of the axial ligands of the heme iron (13). The roman numerals indicate the oxidation state of the heme iron, and His stands for the amino acid residue histidine. The solid and broken arrows indicate rapid and slow reaction steps, respectively. Colour code: black, heme iron with ligands, where the horizontal line through Fe represents a side view of the heme c ring (Figure 9); blue, chemicals added to the protein solution; red, new structural features resulting from the preceding reaction.




Figure 12. Changes with time in the 1D ^1H NMR spectrum of cytochrome c after reducing the ferric cyanide complex with dithionite (220 MHz, 9°C , solvent $^2\text{H}_2\text{O}$).

chrome c contains a methionine side chain of its polypeptide chains in one of the axial heme iron coordination sites. Upon addition of KCN to the protein solution, this methionine is replaced in the binding site on the heme iron by a cyanide ion, and therefore moves out of the heme ring current field. The completion of this reaction can readily be monitored, since different patterns of well-resolved resonance lines are present in the ^1H NMR spectra of ferricytochrome c and its cyanide complex, respectively. After reduction of the heme iron to the ferrous state, the cyanide complex is thermodynamically less stable than the native form of cytochrome c, and in a slow reaction the cyanide in the axial coordination site of the heme iron is again replaced with the natural methionine ligand. The return of the axial methionine into the heme ring current field was monitored in real time (Figure 12) by the appearance of its typical high-field lines in the ^1H NMR spectrum (Figure 10). This experiment resulted in the identification of the nature of this axial ligand in the native, active form of cytochrome c, and further yielded information on the relative thermodynamic stabilities of the four different structures in Figure 11.

As a sideline, comparison of the figures 10 and 12 illustrates the influence of various experimental factors on the appearance of the NMR spectra. The higher resonance frequency and the high temperature used in the experiment of Figure 10, which was recorded around 1980 with Fourier transform spectroscopy, resulted in narrower and more clearly separated peaks in the region 0 to -5 ppm than the experiment in Figure 12, which was recorded in 1968 with continuous-wave NMR spectroscopy at low temperature.

Other early experiments based on the observation of well-resolved resonances in the ^1H NMR spectra of hemoproteins resulted in the characterization of conformation changes during the oxygenation of myoglobin and he-

moglobin (14, 15), and to detailed characterization of the electronic structure of the heme groups in different classes of hemoproteins (16–18).

THE WAY TO NMR STRUCTURES OF PROTEINS

Around 1970 there was a lot of enthusiasm in my laboratory and elsewhere about the success of experiments of the types illustrated by Figures 5 to 12. During the following years, however, there was little further progress toward an NMR method for *de novo* structure determination of proteins. In hindsight this can readily be rationalized: Early successful structural interpretations of NMR data invariably supplemented a previously known low resolution X-ray crystal structure of the same protein. In spite of the high symmetry of the ring current fields (Figure 8) and other sources of local conformation-dependent chemical shifts, including natural paramagnetic centers and extrinsic “paramagnetic shift reagents” (17, 19), the observed chemical shifts could thus in some instances yield exciting new information. In a *de novo* protein structure determination, however, the high symmetry of the local magnetic fields would lead to ambiguities in the structural interpretation of the resulting chemical shifts. Different approaches were therefore called for, and eventually a NMR method for protein structure determination could be based on the following four principal elements:

1. Measurement of NOE upper distance limits as conformational constraints

Nuclear Overhauser effects (NOE) are due to dipolar interactions between different nuclei. The intensity of the NOE is related to the product of the inverse sixth power of the internuclear distance and a correlation function, $f(\tau_c)$, which describes the modulation of the dipole–dipole coupling by stochastic rate processes, with an effective correlation time τ_c :

$$\text{NOE} \propto \frac{1}{\langle r \rangle^6} \cdot f(\tau_c) \quad [1]$$

Although the NOE is a common phenomenon for all combinations of closely spaced nuclear spins, NOEs between pairs of hydrogen atoms are of prime interest for structural studies. A ^1H – ^1H NOE is related to the “through-space” distance between a pair of atoms that are either not at all linked by covalent bonds (intermolecular NOE), or that may be far apart in the amino acid sequence of a polypeptide chain. ^1H – ^1H NOEs can be observed in double-irradiation 1D NMR experiments as the fractional change in intensity of a NMR line that results from preirradiation of another resonance (inset in Figure 13). Although NOE distance measurements had successfully been used for studies of small organic molecules (20), including oligopeptides (21), and observation of NOEs in proteins had also been reported (22, 23), it was not clear whether the desired distance information could actually be obtained in proteins (20, 24, 25). This uncertainty arose because the Brownian motions of large structures in solution are slow, with long effective correla-




Figure 13. NOE build-up curves observed with 1D transient [$^1\text{H}, ^1\text{H}$]-NOE measurements (26). The relative intensities, I_{rel} , of the pre-irradiated resonance line (broken curve) and of the lines experiencing NOEs (solid curves) are plotted *versus* the duration of the mixing period, τ . The experimental scheme used to record transient NOE spectra (inset in the upper right) includes a selective 180° radio-frequency pulse applied to the “pre-irradiated” ^1H NMR line, and a non-selective 90° -pulse applied to the entire ^1H NMR spectrum. The two pulses are separated by the mixing period, τ , and followed by signal acquisition. When working with the crowded spectra of proteins, one records the difference between two spectra obtained with the pre-irradiation $180^\circ(^1\text{H})$ -pulse applied on- and off-resonance, respectively.

tion times, τ_c , for this stochastic process, and proteins contain a dense network of hydrogen atoms. With the combination of these two features, “spin diffusion” could partially or fully deteriorate distance measurements based on ^1H - ^1H NOE experiments (24 – 28).

In the solution NMR experiments discussed here, spin diffusion arises as a consequence of the dependence of the NOE on the inverse sixth power of the internuclear distance, since magnetization transfer between two spins through multiple short steps may be more efficient than a one-step transfer over the longer, direct distance (Figure 14). We used 1D transient NOE difference experiments (inset in Figure 13) (26) and 1D truncated-driven NOE (TOE) difference experiments (27) to record “NOE build-up curves” (Figure 13), which then provided a basis for ^1H - ^1H distance measurements in macromolecules. These studies showed that even more favourable conditions for NOE distance measurements can be found in macromolecules, which have long effective correlation times for the modulation of dipole-dipole couplings, than in low-molecular-weight compounds, for which the condition of “extreme motional narrowing” applies (24, 28).

Analysis of ^1H - ^1H NOE build-up curves recorded with 1D transient NOE experiments (Figure 13) is based on the consideration that during the early phase of the mixing period, direct magnetization transfer from spin 1 to spin 3 (Figure 14) will result in a linear increase of the magnetization on spin 3 with time. In contrast, transfer to atom 3 *via* an intermediate step through atom 2 will have a lag period, even though for long mixing periods, τ , the extent of the magnetization transfer by such “spin diffusion” may for individual pairs of hydrogen atoms exceed that of the direct transfer (Figure 13).




Figure 14. Spin diffusion. Transfer of magnetization between two hydrogen atoms 1 and 3 in the presence of additional hydrogen atoms goes through two competing pathways, *i.e.*, the direct NOE across the distance $r_{1,3}$, and two- or multiple-step spin diffusion *via* intervening hydrogen atoms.

Corresponding considerations apply for the analysis of NOE build-up curves recorded with truncated-driven NOE experiments (27). With proper selection of the duration of the mixing period, one can thus measure highly selective ^1H - ^1H NOEs between distinct pairs of hydrogen atoms, or groups of chemical shift-equivalent hydrogen atoms in proteins in solution.

There is a second criterion that needs to be satisfied for obtaining selective NOEs with 1D NMR experiments, *i.e.*, at least one of the two lines that are connected by the NOE must be sufficiently well resolved (*i.e.*, separated from all other lines in the spectrum) to enable selective radio-frequency pre-irradiation (inset in Figure 13). The previously discussed well-resolved lines in the ^1H NMR spectrum of ferrocycytochrome c (Figure 10) thus had again an important role in enabling us to study the spin physics in the interior of this globular protein with 1D ^1H NMR experiments, as well as to obtain novel structural information. The figure 15 shows a series of highly selective 1D truncated-driven NOE measurements, which were used to determine ^1H NMR assignments for the heme group in a c-type cytochrome (Figure 9) (29).

2. Sequence-specific resonance assignments

Similar to the situation in heme groups (Figure 9), there are closely spaced pairs of hydrogen atoms in neighbouring residues of a polypeptide chain (Figure 16). These can be connected by the observation of “sequential NOEs”. The figure 16 illustrates that NOE-based ^1H NMR assignments for a polypeptide chain can conceptually be considered as a 2-step process. Each amino acid residue represents a “spin system” (4), *i.e.*, it consists of an array of hydrogen atoms including an amide proton (H^{N}), an α -proton (H^{α}), and the side chain protons, which can be connected by steps over three or less covalent bonds through the observation of scalar spin-spin (“through-bond”) couplings (see (4) for the exceptions represented by proline and the aromatic side chains). In contrast, hydrogen atoms located in sequentially neighbouring amino acid residues are separated by at least four covalent bonds. Pairs of neighbouring residues in the sequence can therefore only be connected *via* NOEs manifesting short through-space distances, such as $d_{\alpha\text{N}}$ and d_{NN} . Suitable combinations of intraresidual ^1H - ^1H connectivities established by scalar spin-spin couplings, and inter-residue connectivities established by sequential NOEs enable progressive resonance assignments while “walking




Figure 15. ^1H NMR assignments for the heme group of a c-type cytochrome using ^1H - ^{13}N NOEs (see Figure 7). Bottom trace: 1D ^1H NMR spectrum of ferrocycytochrome c-551 from *Pseudomonas aeruginosa* ($M \approx 12000$). Upper traces: three 1D truncated-driven NOE difference spectra obtained with selective pre-irradiation (indicated by arrows) on the hydrogen atoms β , α and δ (see Figure 9), respectively (360 MHz, 35°C , $^2\text{H}_2\text{O}$ -solution). The NOE peaks are identified with numbers indicating the substituents attached to the corresponding porphyrin ring carbons (Figure 9).

along the polypeptide backbone". In other words, the spin systems of two neighbouring residues can be connected by the intervening H^α - H^N or H^N - H^N sequential NOE connectivities (Figure 16). 1D double-resonance experiments with selective irradiation of the well-resolved amide proton resonances between 8 and 10 ppm in folded BPTI (Figure 5) thus yielded assignments for most of the residues in the regular secondary structure elements (Figure 17) (30). Further assignments were not possible, because the other regions of the polypeptide chain were not represented by well-separated ^1H NMR lines that could have been selectively irradiated in 1D NMR experiments.

3. Two-dimensional (2D) NMR

With the introduction of 2D NMR experiments, and subsequently 3D and 4D NMR experiments, NMR studies of biological macromolecules evolved from intellectually stimulating science to a practical approach for protein structure determination. The foundations of multi-dimensional NMR have been presented in the Nobel lecture by Richard R. Ernst (31). Here, I only want to




Figure 16. Sequential ^1H NMR assignment of proteins. The drawing shows the chemical structure of a -valine-alanine- dipeptide segment in a polypeptide chain. The dotted lines connect groups of hydrogen atoms that are separated by at most three chemical bonds and can therefore be connected using scalar spin-spin couplings. The broken arrows link neighbouring amino acid residues that are separated by short through-space distances, $d_{\alpha\text{N}}$ and d_{NN} , and can therefore be connected by "sequential NOEs".




Figure 17. Sequence-specific resonance assignments for BPTI that were obtained using 1D ^1H NMR experiments (30). The polypeptide backbone is shown, and the assigned residues are identified by indication of hydrogen bonds with their amide protons (the colour code indicates variable exchange rates of the amide protons with the solvent; drawing by Jane Richardson, 1980).

comment on two crucial consequences of multi-dimensional NMR for studies of proteins. First, 2D ^1H NMR enables the recording of selective interactions between pairs of hydrogen atoms, or groups of chemical shift-equivalent hydrogen atoms, without selective irradiation of individual resonance lines. It thus enables a detailed analysis of the entire ^1H NMR spectrum of a protein, which contrasts with the 1D NMR situation of being limited to using only a small number of resolved lines at the periphery of the spectrum. Second, the dispersion of the resonances in a two-dimensional frequency plane affords greatly improved separation of the individual peaks. For example, in Figure 18 the intense lines on the diagonal from the lower left to the upper right correspond to the 1D NMR spectrum, and the weak “cross peaks” in the plane outside of the diagonal manifest selective NOEs between pairs of




Figure 18. Two-dimensional (2D) $[^1\text{H}, ^1\text{H}]$ -NOE spectroscopy ($[^1\text{H}, ^1\text{H}]$ -NOESY). A stacked plot representation of a spectrum of the small protein bull seminal proteinase inhibitor IIA (BUSI IIA, $M \approx 6000$) is shown (500 MHz, 45°C , H_2O -solution).




Figure 19. 2D [$^1\text{H}, ^1\text{H}$]-NOESY spectrum of the plant pathogenesis-related protein P14A ($M \approx 15000$). A contour plot of the spectral region [$\omega_1(^1\text{H}) = 0 - 4.3$ ppm, $\omega_2(^1\text{H}) = 6.3 - 9.5$ ppm] is shown (750 MHz, 30°C , H_2O -solution).




Figure 20. Measurement of NOE build-up curves using 2D [$^1\text{H}, ^1\text{H}$]-NOESY experiments. (A) Six NOESY spectra of BPTI recorded with different mixing times, as indicated on the left. The same 1D cross-section along the ω_1 frequency axis through the diagonal peak of the Phe 33 amide proton is plotted for each spectrum. (B) NOE build-up curves obtained from the analysis of the data in (A); same presentation as in Figure 13, with the relative intensities of the NOESY cross peaks plotted versus the length of the mixing time, τ_m , and the broken line representing the decay of the magnetization on the diagonal peak of the Phe 33 amide proton.




Figure 22. NOESY-COSY connectivity diagram for sequential ^1H NMR assignments using the $d_{\alpha\text{N}}$ sequential NOEs. The data was recorded with BPTI in $^2\text{H}_2\text{O}$ solution. (see text).

4. Structural interpretation of NOE distance constraints

A polypeptide chain with 100 amino acid residues has a length of about 400 Å, whereas NOE-observable distances are shorter than about 5 Å. Observation of a NOE between a pair of hydrogen atoms with assigned chemical shift positions therefore enforces the formation of a ring-like structure (Figure 24). A successful structure determination generates three-dimensional arrangements of the polypeptide chain that simultaneously contain all




Figure 23. Standard presentation of ^1H NMR data leading to sequence-specific resonance assignments and the identification of regular secondary structures in proteins. Experimental data plotted versus the amino acid sequence are shown for the small pheromone protein Er-1 from *Euplotes raikovii* ($M \approx 4500$). $J_{\text{HN}\alpha}$ are scalar spin-spin coupling constants, with small and big values indicated by filled and empty circles, respectively. d_{NN} , $d_{\alpha\text{N}}$ and $d_{\beta\text{N}}$ are distances manifested in sequential NOEs, where strong and weak sequential NOEs are indicated by a thick or a thin line, respectively. Small values of the distances $d_{\alpha\text{N}}(i,i+3)$, $d_{\alpha\beta}(i,i+3)$ and $d_{\alpha\text{N}}(i,i+4)$ are observed by medium-range NOEs linking the given atom types between residues spaced as indicated in the parentheses, and by the short horizontal lines. The locations of three α -helices are indicated at the bottom.




Figure 24. Scheme indicating the relations between an experimental 2D $[^1\text{H}, ^1\text{H}]$ -NOESY spectrum, a polypeptide with the chain ends indicated by N and C, sequence-specific assignments for two hydrogen atoms in the polypeptide chain indicated by circles, and the NOE upper distance constraint derived from the NOESY cross-peak connecting the chemical shift positions of the two assigned hydrogen atoms (see text).

the small and large circular structures imposed by the ensemble of all NOESY cross peaks.

Partial structure determination has been obtained by an empirical approach for the identification of regular secondary structures in polypeptide chains, which relies on recognizing distinct patterns of NOEs (4, 37). For example, in Figure 23 a succession of strong sequential d_{NN} NOEs in combination with the observation of medium-range NOEs in the same polypeptide segment identifies three α -helical structures, which are independently also indicated by successions of small values for the scalar spin-spin couplings $^3J_{\text{HN}\alpha}$ (4, 38).

For the calculation of complete three-dimensional protein structures from NMR data (39), it was quite clear from the outset that an input of quantitative NOE distance measurements would be difficult to obtain. The observed NOEs depend on the proton-proton distance, r , as well as on the effective rotational correlation times, τ_c (Equation [1]). Since for each pair of hydrogen atoms the effective τ_c -value is governed not only by the overall rotational molecular tumbling (Brownian motions), which depends on the size and shape of the protein as well as on the viscosity of the solvent, but can also be affected by intramolecular motions, $f(\tau_c)$ may vary for different pairs of hydrogen atoms in a protein molecule. Additional ambiguities could arise from partial quenching of individual ^1H - ^1H NOEs by competitive spin relaxation processes, for example, spin diffusion (Figure 14), chemical or conformational exchange, and interactions with other nuclear or electronic spins. Furthermore, as a result of rapid intramolecular mobility, a given NOE may be the result of sampling over a range of distances between the two hydrogen atoms of interest (40). In view of these intrinsic limitations for efficient quantitative NOE distance measurements, we decided to use a constant value of the correlation




Figure 25. NMR structure of BUSI IIA (43).

function, $f(\tau_c)$ (Equation [1]), for all ^1H - ^1H combinations in a protein, and to derive only upper limits on the ^1H - ^1H distances from the NOE measurements. In practice, the input for a structure calculation then consists of allowed distance ranges, which are bounded by a NOE upper limit of 3.0 to 5.0 Å, depending on the intensity of the NOE, and a lower limit of 2.0 Å, which represents the sum of the van der Waals radii of the two NOE-connected hydrogen atoms. Although each individual entry in the input data thus has only limited precision, this procedure is robust and can conceptually account for the influence of intramolecular mobility in most of the situations that are commonly expected for the structured parts of globular proteins.

For the initial globular protein structure calculations from NMR data (Figure 25), we used a metric matrix distance geometry algorithm to search for molecular geometries that are consistent with the ensemble of all experimentally determined NOE distance constraints (41–43). Each such calculation ends with the minimization of an error function, and the residual error function value represents a straightforward measure for the success of having found a molecular geometry that satisfies the experimental input data. In view of the aforementioned distance-range format of the input, it is further of keen interest to evaluate the uniqueness of the calculated structure. To this end, the structure calculation is repeated with identical input data but different boundary conditions, and the uniqueness of the resulting NMR structure is judged from the tightness of the fit among the resulting ensemble of conformers. Typically, about 100 conformers are generated, and a sub-group of the 20 conformers with the smallest residual error function values is selected to represent the NMR structure of the protein. The average of the pairwise root-mean-square distances (RMSD) calculated for this bundle of conformers (Figure 1) is then taken as a measure for the precision of the structure determination. Visually, a tight fit of the bundle of conformers indicates regions

where the structure is defined with high precision by the NMR data, whereas structurally disordered polypeptide segments show a large dispersion among the members of the bundle, as exemplified by the two chain ends of the *Antennapedia* homeodomain in Figure 1. In the absence of long-range NOE distance constraints, a properly functioning algorithm for the structure calculation will sample the entire conformation space that is accessible with the given length of the polypeptide chain, as exemplified by the “unstructured” tail of the bovine prion protein in Figure 4.

NMR IN STRUCTURAL BIOLOGY

Standard protocol for NMR structure determination of biological macromolecules

The protocol for NMR structure determination includes the preparation of a homogeneous protein solution, the recording and handling of the NMR data sets, and the structural interpretation of the NMR data (Table 1). The techniques used in 1984 for the structure determination of bull seminal proteinase inhibitor IIA are listed in Table 1; the four steps of the structural interpretation (III in Table 1) were performed separately, although the result of the first round of constraint collection and structure calculation was subsequently used for additional checks on the sequence-specific resonance assignments as well as on the collection of conformational constraints. Since 1984, the protocol outlined in Table 1 has been used in over 3'000 NMR structure determinations of proteins and nucleic acids, and greatly improved experimental techniques have been incorporated into this general scheme.

Major advances in the experimental techniques for NMR structure determination were spurred by the introduction of methods for the production of recombinant proteins labelled with stable isotopes, in particular

Table 1. Standard protocol for NMR structure determination of proteins

Step ^a	BUSI IIA ^b
I Sample preparation	Protein isolated from natural source; natural isotope distribution; 16 mM solutions in H ₂ O and in ² H ₂ O, respectively
II NMR spectroscopy	2D ¹ H NMR
IIIa Resonance assignments	Sequential NOEs
IIIb Conformational constraints	[¹ H, ¹ H]-NOEs, ³ J _{HNaα'}, ³J_{α\beta}}
IIIc Structure calculation	Metric matrix distance geometry
IIId Structure refinement	Restrained energy minimization

^a The structural interpretation of the NMR data, III, is somewhat arbitrarily divided up into four steps; in practice, one goes through multiple cycles of collection of conformational constraints (IIIb) and structure calculation (IIIc), and the completion of the sequence-specific assignments (IIIa) as well as the structure refinement (IIId) may also be part of this iterative approach.

^b This column lists the techniques used in the first structure determination of a globular protein in 1984 (43).




Figure 26. Three-dimensional (3D) ^{15}N -resolved $[^1\text{H}, ^1\text{H}]$ -NOESY spectrum (600 MHz, 28°C , H_2O -solution) of the DNA-binding domain of the P22 c2 repressor ($M \approx 10000$, uniformly ^{15}N -labeled).

carbon-13, nitrogen-15 and deuterium (44, 45). For example, this opened the way for efficient use of heteronuclear NMR techniques with proteins, such as 3D $[^1\text{H}, ^{13}\text{C}, ^{15}\text{N}]$ -triple resonance experiments, 3D ^{13}C - or ^{15}N -resolved $[^1\text{H}, ^1\text{H}]$ -NOESY (Figure 26) (46, 47), and the use of heteronuclear filters (48). Important advances have also been made with the methods of structure calculation, where the c.p.u. time needed for the calculation of a small protein structure has been reduced from about one day in 1984 (41, 42) to a few seconds (49). Currently, intense work is focused on the automation and combined execution of the individual steps in the structural interpretation of the NMR data (50, 51). It is beyond the scope of this article to present in detail the wide range of beautiful novel experimental approaches developed by the community of macromolecular NMR spectroscopists during the past 15 years, which now enable studies with ever more complex systems (Figures 1–4).

Globular protein structures in solution

The static picture of a protein molecule obtained from the standard protocol for structure determination (Table 1) typically shows variable precision of the structure determination along the polypeptide chain, as manifested by the variations in the closeness of the fit among the bundle of conformers used to represent the NMR structure (Figure 1). Even in proteins where the entire polypeptide chain is part of the global fold, increased disorder is observed toward the periphery of the surface side chains (Figure 27). This pronounced surface disorder, which typically also involves the ends of the polypeptide chain, is in most instances the only significant difference between corresponding globular protein structures in single crystals and in solution (5). In part the increased disorder in solution arises because of a scarcity of packing constraints, when compared with the protein core, and a concomitant scarcity of NOE distance constraints. With the additional use of NMR spin relaxation measurements (24, 28, 33, 52), one can distinguish between static disorder, which would then presumably arise from the scarcity of constraints,




Figure 27. NMR structure of BPTI represented by a bundle of 20 conformers superimposed for best fit of the polypeptide backbone. The polypeptide backbone is green, core side-chains are blue, and solvent-accessible surface side-chains are red.

and dynamic disorder, with intramolecular motions on the nanosecond and sub-nanosecond time scale. Overall, quite independent of the dynamics issue, the observation of partially folded polypeptide chains in solution (Figures 1, 4 and 27) is important complementary information to the data that can be obtained by studies in crystals. It is also the main reason why the quality of a NMR structure determination is not usually characterized by a single, global parameter (4–6).

An important extension of the characterization of proteins in solution resulted from high resolution NMR studies of protein hydration. Thereby the location of hydration waters is determined by the observation of NOEs between water protons and hydrogen atoms of the polypeptide chain (53). Because of the dependence of the NOE on the inverse sixth power of the ^1H – ^1H distance, only one layer of hydration water molecules is observed (Figure 28). For the hydration studies, the dependence of the NOE intensity




Figure 28. Molecular model of hydrated BPTI in H_2O solution. The drawing shows an all-heavy-atom presentation of one of the conformers in Figure 27 (yellow) covered with a layer of hydration water molecules (dotted blue spheres).




Figure 29. Intramolecular rate processes in BPTI. The polypeptide backbone is represented by a grey spline function through the α -carbon positions, with the thickness of the line representing the spread of the bundle of 20 conformers in Figure 27. Individual rate processes and their frequencies are indicated with the following colour code: dark blue, magenta and red: ring flips of phenylalanine and tyrosine; yellow, exchange between the R- and S-chiral forms of the disulfide bond at the top; cyan, interior hydration water molecules, with indication of the exchange rates with the bulk water.

on the correlation function describing the stochastic modulation of the dipole–dipole coupling between the interacting protons (Equation [1]) has a key role. The value of $f(\tau_c)$ may be governed either by the Brownian rotational tumbling of the hydrated protein molecule, or by interruption of the dipolar interaction through translational diffusion of the water molecules relative to the protein surface, whichever is faster. On this basis it could be established that surface hydration of peptides and proteins is characterized by very short residence times of the water molecules in the hydration sites, in the range from about 20 to 300 picoseconds at 10°C. This result presents an intuitive rationale for the generally observed dynamic disorder of the protein surface structure (Figure 27), and indicates that the dehydration of the polypeptide surface will hardly ever be a rate-limiting step either in protein folding or in intermolecular interactions with proteins.

The BPTI crystal structure contains four interior hydration water molecules. These are an integral part of the molecular architecture and are inaccessible to the solvent in a rigid model of the three-dimensional structure (Figure 29). The chemical shift dispersion anticipated for these four water molecules on the basis of the considerations in Figures 5–8 was not observed. This degeneracy of the chemical shifts of bound water and bulk water was found to be due to rapid exchange of water molecules in and out of the protein molecule, with an upper limit on the life-time of about 1 millisecond (53; as indicated in Figure 29 by the number in parentheses, the actual life times

for individual waters may be significantly shorter). Rapid exchange of interior hydration water molecules appears to be a general property of globular proteins, and was also observed for water molecules located in protein–DNA interfaces, for example, in the DNA complex with the *Antennapedia* homeodomain (Figure 2).

Another intriguing NMR observation of internal protein mobility are the 180°-ring flipping motions of phenylalanine and tyrosine (54). Observation of these “ring flips” on the millisecond to microsecond time scale (Figure 29) was a genuine surprise for the following reasons: in the refined X-ray crystal structure of BPTI reported in 1975, the aromatic rings of phenylalanine and tyrosine are among the best-defined side chains, with the smallest temperature factors. In each ring the relative values of the temperature factors for the individual atoms increase toward the periphery, so that the largest positional uncertainty is indicated for the peripheral ζ -carbon atom on the symmetry axis through the C^β – C^γ bond, rather than for the four δ - and ϵ -ring carbon atoms, which undergo extensive movements during the ring flips. Theoretical studies then resolved this apparent contradiction: the crystallographic temperature factors sample multiple rotation states about the C^α – C^β bond, but they do not manifest the ring flips because the populations of all non-equilibrium rotational states about the C^β – C^γ bond are vanishingly small. Although the flipping motions about the C^β – C^γ bond have low frequencies (Figure 29), they are very rapid 180°-rotations connecting two indistinguishable equilibrium orientations of the ring. Similar to the exchange of internal hydration waters, the ring flip phenomenon is a general feature of globular proteins, manifesting ubiquitous low-frequency internal motions that have activation energies of 60 to 100 kJmol⁻¹, amplitudes larger than 1 Å, activation volumes of about 50 Å³, and involve concerted displacement of numerous groups of atoms. Combined with sequence-specific NMR assignments, these experiments provide high spatio-temporal resolution for the description of rate processes in proteins. In Figure 29 this is illustrated by a mapping of the frequencies for ring flips and water exchange onto the NMR structure of BPTI.

In addition to the ring flips and the exchange of internal hydration water molecules, the figure 29 includes data on the exchange of a disulfide bond between the R and S chiral states. In contrast to the other two phenomena, this rate process connects two different molecular structures.

Although all the data collected in Figure 29 have been known for more than a decade, and some of them for nearly three decades, no widely accepted functional interpretation of these low-frequency motional processes has been advanced. The same holds for the conformational equilibria manifested by the “protection factors” governing the amide proton exchange rates in folded proteins (Figures 17 and 21). Quite possibly these NMR measurements are ahead of their times, and represent a source for future novel insights into structure–function correlations in proteins.




Figure 30. NMR structure determination and molecular weight. The horizontal axis covers the molecular weight-range 0 – 110'000. The left side shows the molecular weight distribution of the NMR structures in the protein databank (December 2000). On the right, the structure of the protein OmpX from *E. coli* ($M \approx 18000$) in DHPC micelles ($M \approx 70\ 000$ for the mixed micelles) indicates along the horizontal axis the approximate molecular weight range for NMR structure determination of mixed micelles with membrane proteins that is presently accessible with the NMR techniques of Figures 31 – 33.

OUTLOOK TO NMR APPLICATIONS IN STRUCTURAL AND FUNCTIONAL PROTEOMICS

With the availability of a rapidly increasing number of completely sequenced genomes, new challenges arise for the methods used for three-dimensional structure determination. On the one hand, “structural genomics” initiatives in several leading research centers focus on the development of technology for high-throughput structure determination to generate a comprehensive atlas of protein folds, so that remaining gaps could be filled by structure prediction methods. There is clearly a lot of room to further enhance the efficiency of each step of the NMR structure determination procedure (Table 1). On the other hand, we face the situation that newly determined protein structures should enable us to predict novel functions, whereas in classical structural biology one encounters more typically the challenge of rationalizing known functions on the basis of the three-dimensional structure. This section describes some recent work in our laboratory that may eventually contribute to future strategies for the discovery of new physiological functions from molecular structure data.

It has been widely recognized that supplementing the determination of new protein folds with data on intermolecular interactions may provide a key for the identification of unknown gene functions. Since efficient use of conventional NMR spectroscopy in solution had been limited to particle sizes with molecular weights up to about 30'000 (Figure 30, left side), a new chal-




Figure 31. Transverse relaxation-optimized [^{15}N , ^1H]-correlation spectroscopy (TROSY). A spectrum of the uniformly ^2H - and ^{15}N -labeled membrane protein OmpX reconstituted in DHPC micelles is shown (750 MHz, 30°C, H_2O -solution).

lenge for solution NMR techniques then arose from the fact that the supramolecular structures resulting from interactions of two or several proteins, or of other macromolecular components tend to have high molecular weights. Although a 30'000-Dalton size limit allowed to work with a large pool of physiologically interesting proteins, it was hardly compatible with extensive use of NMR for studies of such supramolecular structures. For example, this size limit would severely narrow down the range of potential receptor systems accessible to NMR in drug discovery projects (55, 56), restrict studies of protein–nucleic acid complexes (Figure 2) to a small number of systems with modest size, and prevent the use of solution NMR for studies of membrane proteins, since these have to be reconstituted and solubilized in mixed micelles with detergents or lipids (Figure 30, right side). A few years ago this limitation was successfully challenged, since the size-range for applications of solution NMR techniques could be significantly extended through the introduction of transverse relaxation-optimized spectroscopy (TROSY) (57). As an illustration, the figure 31 shows a [^{15}N , ^1H]-TROSY correlation spectrum of a membrane protein reconstituted in detergent micelles. Sharp, well-separated peaks are obtained in spite of the large size of the mixed micelles (Figure 30, right side). With the use of the TROSY-principle, the more complex NMR experiments needed for a structure determination with [^2H , ^{13}C , ^{15}N]-labelled proteins can also be obtained with high quality (58), so that solution NMR can now be used for *de novo* membrane protein structure determination (59).

The improved quality of the NMR spectra of large structures with the use of TROSY can be qualitatively rationalized by the following considerations. As has previously been indicated in the discussion on NOEs (Equation [1]), the appearance of solution NMR spectra is intimately related to the effective cor-




Figure 32. Schematic drawing of the free induction decay (FID) for a large protein with and without the use of TROSY. The period covered along the time axis is of the order of 100 milliseconds. The FID represents the primary recording of the NMR data in the “time domain”, from which the frequency domain spectrum (for example, Figures 18, 19 and 31) is obtained by Fourier transformation (31, 33).

relation time, τ_c , which characterizes the thermal motions of the molecule considered. The increase of the τ_c -values in larger structures results in line broadening due to rapid transverse spin-relaxation. For example, if the spectrum of Figure 31 had been recorded with conventional NMR techniques, most of the resonance lines would not be individually resolved due to severe line broadening, and one would have experienced a severe loss of sensitivity for detection of the NMR signals. The reduced sensitivity can be readily appreciated by examination of the “free induction decays” (FID) in the time domain data. The figure 32 shows that rapid loss of magnetization in a conventional NMR experiment with a large structure can be slowed down by the use of TROSY, which corresponds to the reduction of the line width in the frequency domain spectrum. The impact on the sensitivity is visualized in Figure 33 with a simplified scheme for a 2D correlation experiment, which includes two magnetization transfer periods, and the evolution and acquisition periods. During evolution and acquisition, the system is not subjected to external perturbations, and loss of magnetization occurs at a rate determined by the “transverse relaxation time”. It is readily apparent that rapid loss of magnetization in a conventional NMR experiment (upper trace in Figure 32) leads to weak or even vanishing signal intensity at the end of the evolution period, and accordingly the sensitivity for detection of the signal during the acquisition period is very low. In contrast, one can obtain much improved sensitivity with the use of TROSY (lower trace in Figure 32), since plenty of signal intensity will be preserved at the end of the evolution period, and the signal can be recorded with high signal-to-noise ratio during a large portion of the ac-




Figure 33. Basic features of a 2D [^{15}N , ^1H]-correlation experiment. On the left, ^1H and ^{15}N indicate radio-frequency channels for irradiation of these isotopes. Yellow shading identifies the $^1\text{H} \rightarrow ^{15}\text{N}$ and $^{15}\text{N} \rightarrow ^1\text{H}$ magnetization transfer periods, T , and pink shading the evolution and acquisition periods, t_1 and t_2 . During t_1 and t_2 the decay of the magnetization (FID) is schematically indicated. The overall duration of the experiment, $2T+t_1+t_2$, can be of the order of 10 to 1000 milliseconds, depending on the size of the structure studied and the intended purpose of the measurement.

quisition period. Following such considerations for minimizing the loss of magnetization during all four time periods indicated in Figure 33, and with additional optimization of the magnetization transfer techniques (60), solution NMR spectra have by now been recorded for structures with molecular weights up to 870'000. In the spectrum of the co-chaperonin GroES bound to the chaperonin GroEL (Figure 34), resonance lines that provided novel in-




Figure 34. [^{15}N , ^1H]-correlation spectrum of the co-chaperonin GroES ($M \approx 70000$) bound to the chaperonin GroEL ($M \approx 800000$). The spectrum was recorded with a [^{15}N , ^1H]-CRIPT-TROSY experiment (60, 61). The red and blue colouring is explained in the text.




Figure 35. NMR structure of the membrane protein OmpX reconstituted in DHPC micelles. Front and back views of a space-filling model of OmpX are shown, with pink and green coloring identifying the surface area of the protein that is in contact with DHPC molecules in the mixed micelles. The two broken horizontal lines indicate the thickness of the lipid phase in the *E. coli* cell membrane.

formation on structural and dynamic features of the GroES–GroEL interface are colored in red and blue (61).

The principal contributions to the rate of transverse spin-relaxation can be traced to two different types of interactions, *i.e.*, dipole–dipole coupling of the observed spin with other, nearby spins, and chemical shift anisotropy (CSA) (28). These interactions are modulated by the stochastic rotational motions in solution, and as a consequence the rate of transverse relaxation increases for larger structures with slower Brownian motions. TROSY exploits constructive interference between the two relaxation mechanisms, and actually uses CSA-relaxation at high fields to cancel the dipolar relaxation (57). In this way the appearance of the NMR spectrum is effectively uncoupled from the Brownian motions, which then enables the recording of solution NMR spectra with large structures.

It remains to be seen how these new NMR techniques will be employed most profitably in the future. Intriguing possibilities include that NMR can now be employed in drug discovery projects with very large receptors (55, 56). Combined with suitable isotope-labelling strategies, TROSY-based NMR techniques have also been shown to provide a powerful approach for investigations of intermolecular interactions in supramolecular structures with two or several macromolecular components (62, 63). In these applications, a detailed structural interpretation of the NMR data will in most instances be dependent on the availability of an independently determined atomic-resolution structure for one or multiple components, which may have been obtained either by NMR in solution or by diffraction methods in single crys-

Students, Research associates, Technical staff 1970–2002

Bernar Hindermann	Klaus Löffl	Fay Sroog	Daniel Schmitter	Barthold von Freyberg	Gerhard Frenk	Jeanette Pison
Jean-Paul Meislin	Kristofl Nagypatai	Dominique Marlot	Manuel Günter	Christoph Lechner	Walter Sailer	Lutz Grottel
André Messon	Stephen J. Parkley	Stefan H. P. Zuberhöp	Dezar D. Radtke	Johanna Müller	Konstantin Parnasch	Rajkumar Vohra
Regula M. Keller	Sammy L. Gordon	Fransiska Durrmussan	Francesco Riva	Miguel B. Gil	Ramona Hug	Reto Honek
Jay Gals	Lothar Gier	Arthur Fanti	Hydroly Hargreaves	John P. O'Donnell	Cyril Ansel	Songyan Lee
Kudolf Bäumann	Andreas Galle	Kenneth R. Meis	Daniel Baumann	Barbara Lelling	Roland Riek	Nilsu Akalinçli
Bernard Dorjat	David Platt	Tammy Havel	Bernard Th. Farnes II	John H. Sauerbrey	Francisco Lopez-Garcia	Christiane A. Lysek
Jill Hochmann	Werner A. Bruch	Michael P. Strickland	Christopher Weber	Toshiko Maruta	Marino Moro	Sayma Banger
Gregory Grefenstätt	Heverich Huber	Paula V. Wroblek	Hao Guojun	Marin Brivic	Prud'hu	Christina Heff
Klaus Eder	Albert Siggler	Matthias Hecker	Peter Gütler	Walteri Anstich-Berth	Hail Dasso	Shirley Hainey
Robert Hentsch	Gerdhard Weber	Daniel Neuhäuser	FRANCESCO ARRE	Peter Lugliński	AKIHIKO UO	ROBERT L. J. Koller
Gabriele Kurr	Christoph Wittler	Peter Schuler	Tatjana Ma	Christina Schütz	Michael Salemann	Olga Kruza
Joana M. Targovska	Clark Browne	Sara Mäkelä	Victor Göttsch	Walteri Anstich-Berth	Alexander G. Sobol	Christina Heff
Andreas Göbel	Julia Chavesmayh	WALTER LUGLI	Barbara Messaote	Siggi Morigo	Fred F. Dandinger	Seal Vogel
Antonella Schumacher	Marlene Meier	Selma Kälin	Greg D. Eccles	Jan Johansson	Chih-yuen Ning	Touraj Ehsanizadeh
Kurt Gasser	Sudhanya Ramaswamy	Sara Mamed	Yunqin Chen	Gregg Singh	Magdalene Böckli	Christina Schumacher
Richard Wagner	Hubertus Göl	Walter J. Chazin	Doris Kari	Oliver Schenk	Yves-Laurent Vieille	Francesca Florio
Arno Suter	Mike Göl	Michael H. Frey	Herve Durrbin	Niklaus Schwefel	Ralph Zinn	Patrick Battersdorf
Mauro De Benedicis	Johanna Schaffner	Wolfgang Gork	Hugo Amacker	Gabriel Hänggi	Michael Hochuli	Simona Hommann
Peter Sauer	Markus Schneider	Sven Nykjaer	Wolfgang Hämli	Ram Kowal	Britt-Marie Tobi	Vivian Salvo Noya
Andreas J. Gillies	Hans Sauer	Sudhinda Ghosh	Ivan Weissauer	Helen Kowal	Oliver Lutz	Valérie Fasel
Miguel Loria	Wayne E. Blumenthal	Alan D. Kline	Kristina V. H. Chaly	Marcel Origer	Jörg Pieper	Hart A. Bähler
Hans-Ulrich Grewerich	Andi Ruppel	Alexander N. Lubchenco	Daniel Bruner	Ralph Durrbin	Tien Y. Tsang	Helmut Teller
Walter Müller	Yvonne Havelier	Christel Myerwald	Rolf Meyer	Matthias Gohl	Thorsten Lutz	Ahmed Ghomri
Larry K. Brown	Selma Hüter	CHANG MYERWALD	Frank Sauer	SERGE ARRA	Christine von Salzenstein	Laura Nover
Rolf Honek	Fredrick Spack	Thomas Schumann	Marie Dos	Maria Pia Barilla	Roberto Fattorusso	Subodh Kumar
Alexandra E. Frey	Olga Takemura	FRANCIS STREIB	Agnieszka Fieda	Oliver Farnoch-Estratin	Hans-Joachim Gruber	Ida Ares
CHRISTOPH BLOCH	MARIN BRIVIC	Stefan Wang	James Verdell	Carsten Mueselbauer	Richard Wimmer	Yunqin Chen
Vivian Salvo	Stefan Wipfler	Norbert Müller	Thomas A. Gajarski	Oliver Zerbe	Martha Gohl	Daniel Pflanz
Laszlo Szekely-Gasz	Shobhana Murali	Luigi Jenny	Karl Bernell	Birchitta Pfisterer	Barbara Zangger	Martina Wipfler
Jürgen Lubbenstein	Samuelina Hauer	Sergio Galle	Oliver Spethmann	Helen Höl	Turker Homann	Isabelle Allen
Peter Sauerbrey	Alexandre Ametico	Rolf Götter	Joséphine M. Moos	Oliver Chwastick	Kai Schwan	Christine Anst

Figure 36. My collaborators at the ETH Zürich from 1970 to 2002.

tals. Applications of the new NMR techniques for *de novo* determination of large structures appears to be particularly attractive for, but not limited to, nucleic acid–protein complexes and small membrane proteins reconstituted in soluble detergent or lipid micelles. The characterization of a membrane protein structure has also been extended to include the detergents in the mixed micelles (Figure 35) (64), since the areas of the protein surface in contact with the detergent molecules could be delineated by the observation of intermolecular NOEs between hydrogen atoms of the protein and the detergent. In these particular mixed micelles this surface area is virtually identical to the protein surface in contact with the lipid phase in the biological membrane. Since this leaves both ends of the β -barrel of OmpX freely accessible to the aqueous solvent, this reconstitution system should also be suitable for functional studies of the membrane protein with NMR.

This last section of my presentation certainly indicates only a narrow range of potential further developments of solution NMR techniques and their applications in structural biology and structural proteomics. I look forward with great expectations to the future evolution of this awesome and beautiful technique, which has given me so many years of joy and excitement in studies of the molecules of life.

ACKNOWLEDGMENTS

From 1970 through 2002, 229 students, postdoctoral research associates, and technical and administrative staff worked with me at the ETH Zürich. I am deeply indebted to all of them for their enthusiasm and dedication, which resulted in the work summarized in this article. Their names are collected in Figure 36, and individual contributions can be recognized in the reference list. Although human minds stand behind all progress in science, the success

of our research projects also depended critically on financial resources. I would like to acknowledge the ETH Zürich, the Swiss National Science Foundation, the Kommission für Technologie und Innovation (KTI), Bruker-Biospin AG, and the Scripps Research Institute in La Jolla, CA, USA, for their support.

REFERENCES

1. Qian, Y.Q., Billeter, M., Otting, G., Müller, M., Gehring, W.J. and Wüthrich, K. (1989) *Cell* **59**, 573–580. The structure of the *Antennapedia* homeodomain determined by NMR spectroscopy in solution: comparison with prokaryotic repressors.
2. Otting, G., Qian, Y.Q., Billeter, M., Müller, M., Affolter, M., Gehring, W.J. and Wüthrich, K. (1990) *EMBO J.* **9**, 3085–3092. Protein–DNA contacts in the structure of a homeodomain–DNA complex determined by nuclear magnetic resonance spectroscopy in solution.
3. Lopez Garcia, F., Zahn, R., Riek, R. and Wüthrich, K. (2000) *Proc. Natl. Acad. Sci. USA* **97**, 8334–8399. NMR structure of the bovine prion protein.
4. Wüthrich, K. (1986) *NMR of Proteins and Nucleic Acids*, Wiley, New York.
5. Wüthrich, K. (1995) *Acta Cryst. D* **51**, 249–270. NMR—this other method for protein and nucleic acid structure determination.
6. Wüthrich, K. (1995) *NMR in Structural Biology: A Collection of Papers by Kurt Wüthrich*, World Scientific, Singapore.
7. Wüthrich, K. (2001) *Nature Struct. Biol.* **8**, 923–925. The way to NMR structures of proteins.
8. Saunders, M. and Wishnia, A. (1958) *Ann. N.Y. Acad. Sci.* **70**, 870–874. Nuclear magnetic resonance spectra of proteins.
9. Jardetzky, O. and Jardetzky, C.D. (1958) *J. Biol. Chem.* **233**, 383–387. Proton magnetic resonance spectra of amino acids.
10. Kowalsky, A. (1962) *J. Biol. Chem.* **237**, 1807–1819. Nuclear magnetic resonance studies of proteins.
11. Mandel, M. (1965) *J. Biol. Chem.* **240**, 1586–1592. Proton magnetic resonance spectra of some proteins.
12. McDonald, C.C. and Phillips, W.D. (1967) *J. Amer. Chem. Soc.* **89**, 6332–6341. Manifestations of the tertiary structures of proteins in high-frequency nuclear magnetic resonance.
13. Wüthrich, K. (1969) *Proc. Natl. Acad. Sci. USA* **63**, 1071–1078. High resolution proton nuclear magnetic resonance spectroscopy of cytochrome c.
14. Shulman, R.G., Ogawa, S., Wüthrich, K., Yamane, T., Peisach, J. and Blumberg, W.E. (1969) *Science* **165**, 251–257. The absence of “heme–heme” interactions in hemoglobin.
15. Davis, D.G., Lindstrom, T.R., Mock, N.H., Baldassare, J.J., Charache, S., Jones, R.T. and Ho, C. (1971) *J. Mol. Biol.* **60**, 101–111. NMR studies of hemoglobins VI: heme proton spectra of human deoxyhemoglobins and their relevance to the nature of cooperative oxygenation of haemoglobin.
16. Kurland, R.J., Davis, D.G. and Ho, C. (1968) *J. Amer. Chem. Soc.* **90**, 2700–2701. Paramagnetic proton NMR shifts of metmyoglobin, methemoglobin, and hemin derivatives.
17. Wüthrich, K. (1970) *Structure and Bonding* **8**, 53–121. Structural studies of hemes and hemoproteins by nuclear magnetic resonance spectroscopy.
18. Shulman, R.G., Glarum, S.H. and Karplus, M. (1971) *J. Mol. Biol.* **57**, 93–115. Electronic structure of cyanide complexes of hemes and heme proteins.
19. Wüthrich, K. (1976) *NMR in Biological Research: Peptides and Proteins*, North Holland, Amsterdam.

20. Noggle, J.H. and Schirmer, R.E. (1971) *The Nuclear Overhauser Effect*, Academic Press, New York.
21. Gibbons, W.A., Alms, H., Bockman, R.S. and Wyssbrod, H.R. (1972) *Biochemistry* **11**, 1721–1725. Homonuclear INDORE spectroscopy as a means of simplifying and analyzing proton magnetic resonance spectra of peptides and as a basis for determining secondary and tertiary conformations of complex peptides.
22. Gupta, R.K. and Redfield, A.G. (1970) *Science* **169**, 1204–1206. Double resonance NMR observation of electron exchange between ferri- and ferrocyclochrome *c*.
23. Balaram, P., Bothner-By, A.A. and Dadok, J. (1972) *J. Amer. Chem. Soc.* **94**, 4015–4017. Negative nuclear Overhauser effects as probes of macromolecular structure.
24. Solomon, I. (1955) *Phys. Rev.* **99**, 559–565. Relaxation processes in a system of two spins.
25. Kalk, A. and Berendsen, H.J.C. (1976) *J. Magn. Reson.* **24**, 343–366. Proton magnetic relaxation and spin diffusion in proteins.
26. Gordon, S.L. and Wüthrich, K. (1978) *J. Amer. Chem. Soc.* **100**, 7094–7096. Transient proton–proton Overhauser effects in horse ferrocyclochrome *c*.
27. Wagner, G. and Wüthrich, K. (1979) *J. Magn. Reson.* **33**, 675–680. Truncated driven nuclear Overhauser effect (TOE): a new technique for studies of selective ¹H–¹H Overhauser effects in the presence of spin diffusion.
28. Abragam, A. (1961) *Principles of Nuclear Magnetism*, Clarendon Press, Oxford.
29. Keller, R.M. and Wüthrich, K. (1978) *Biochem. Biophys. Res. Comm.* **83**, 1132–1139. Evolutionary change of the heme *c* electronic structure: ferricytochrome *c*-551 from *Pseudomonas aeruginosa* and horse heart ferricytochrome *c*.
30. Dubs, A., Wagner, G. and Wüthrich, K. (1979) *Biochim. Biophys. Acta* **577**, 177–194. Individual assignments of amide proton resonances in the proton NMR spectrum of the basic pancreatic trypsin inhibitor.
31. Ernst, R.R. (1992) *Angew. Chemie –Int. Ed.* **31**, 805–823. Nuclear magnetic resonance Fourier transform spectroscopy.
32. Wider, G., Macura, S., Anil Kumar, Ernst, R.R. and Wüthrich, K. (1984) *J. Magn. Reson.* **56**, 207–234. Homonuclear two-dimensional ¹H NMR of proteins: experimental procedures.
33. Ernst, R.R., Bodenhausen, G. and Wokaun, A. (1987) *Principles of Nuclear Magnetic Resonance in One and Two Dimensions*, Oxford University Press, Oxford.
34. Wagner, G. and Wüthrich, K. (1982) *J. Mol. Biol.* **155**, 347–366. Sequential resonance assignments in protein ¹H nuclear magnetic resonance spectra: basic pancreatic trypsin inhibitor.
35. Wagner, G., Anil-Kumar and Wüthrich, K. (1981) *Eur. J. Biochem.* **114**, 375–384. Systematic application of two-dimensional ¹H nuclear magnetic resonance techniques for studies of proteins 2: combined use of correlated spectroscopy and nuclear Overhauser spectroscopy for sequential assignments of backbone resonances and elucidation of polypeptide secondary structures.
36. Billeter, M., Braun, W. and Wüthrich, K. (1982) *J. Mol. Biol.* **155**, 321–346. Sequential resonance assignments in protein ¹H nuclear magnetic resonance spectra: computation of sterically allowed proton–proton distances and statistical analysis of proton–proton distances in single crystal protein conformations.
37. Wüthrich, K., Billeter, M. and Braun, W. (1984) *J. Mol. Biol.* **180**, 715–740. Polypeptide secondary structure determination by nuclear magnetic resonance observation of short proton–proton distances.
38. Pardi, A., Billeter, M. and Wüthrich, K. (1984) *J. Mol. Biol.* **180**, 741–751. Calibration of the angular dependence of the amide proton–C^α proton coupling constants, ³J_{HNa}, in a globular protein: use of ³J_{HNa} for identification of helical secondary structure.
39. Wüthrich, K., Wider, G., Wagner, G. and Braun, W. (1982) *J. Mol. Biol.* **155**, 311–319. Sequential resonance assignments as a basis for determination of spatial protein structures by high resolution proton nuclear magnetic resonance.

40. Braun, W., Bösch, C., Brown, L.R., Gö, N. and Wüthrich, K. (1981) *Biochim. Biophys. Acta* **667**, 377–396. Combined use of proton–proton Overhauser enhancements and a distance geometry algorithm for determination of polypeptide conformations: application to micelle-bound glucagon.
41. Havel, T.F. and Wüthrich, K. (1984) *Bull. Math. Biol.* **46**, 673–698. A distance geometry program for determining the structures of small proteins and other macromolecules from nuclear magnetic resonance measurements of intramolecular ^1H – ^1H proximities in solution.
42. Havel, T.F. and Wüthrich, K. (1985) *J. Mol. Biol.* **182**, 281–294. An evaluation of the combined use of nuclear magnetic resonance and distance geometry for the determination of protein conformations in solution.
43. Williamson, M.P., Havel, T.F. and Wüthrich, K. (1985) *J. Mol. Biol.* **182**, 295–315. Solution conformation of proteinase inhibitor IIA from bull seminal plasma by ^1H nuclear magnetic resonance and distance geometry.
44. Wagner, G. (1993) *J. Biomol. NMR* **3**, 375–385. Prospects for NMR of large proteins.
45. Kay, L.E. and Gardner, K.H. (1997) *Curr. Opin. Struct. Biol.* **7**, 564–570. Solution NMR spectroscopy beyond 25 kDa.
46. Bax, A. and Grzesiek, S. (1993) *Accounts Chem. Res.* **26**, 131–138. Methodological advances in protein NMR.
47. Wider, G. (1998) *Prog. Nucl. Magn. Reson. Spectr.* **32**, 193–275. Technical aspects of NMR spectroscopy with biological macromolecules and studies of hydration in solution.
48. Otting, G. and Wüthrich, K. (1990) *Q. Rev. Biophys.* **23**, 39–96. Heteronuclear filters in two-dimensional [^1H , ^1H]-NMR spectroscopy: combined use with isotope labelling for studies of macromolecular conformation and intermolecular interactions.
49. Güntert, P. (1998) *Q. Rev. Biophys.* **31**, 145–237. Structure calculation of biological macromolecules from NMR data.
50. Moseley, H.N.B. & Montelione, G.T. (1999) *Curr. Opin. Struct. Biol.* **9**, 635–642. Automated analysis of NMR assignments and structures for proteins.
51. Herrmann, T., Güntert, P. and Wüthrich, K. (2002) *J. Biomol. NMR* **24**, 171–189. Protein NMR structure determination with automated NOE-identification in the NOESY spectra using the new software ATNOS.
52. Luginbühl, P. and Wüthrich, K. (2002) *Progr. Nucl. Magn. Reson. Spectr.* **40**, 199–247. Semi-classical nuclear spin relaxation theory revisited for use with biological macromolecules.
53. Otting, G., Liepinsh, E. and Wüthrich, K. (1991) *Science* **254**, 974–980. Protein hydration in aqueous solution.
54. Wüthrich, K. and Wagner, G. (1975) *FEBS Lett.* **50**, 265–268. NMR investigations of the dynamics of the aromatic amino acid residues in the basic pancreatic trypsin inhibitor.
55. Shuker S.B., Hajduk P.J., Meadows R.P. and Fesik S.W. (1996) *Science* **274**, 1531–1534. Discovering high-affinity ligands for proteins: SAR by NMR.
56. Pellecchia, M., Sem, D.S. and Wüthrich, K. (2002) *Nature Rev. Drug Disc.* **1**, 211–219. NMR in drug discovery.
57. Pervushin, K., Riek, R., Wider, G. and Wüthrich, K. (1997) *Proc. Natl. Acad. Sci. USA* **94**, 12366–12371. Attenuated T_2 relaxation by mutual cancellation of dipole–dipole coupling and chemical shift anisotropy indicates an avenue to NMR structures of very large biological macromolecules in solution.
58. Salzmann, M., Pervushin, K., Wider, G., Senn, H. and Wüthrich, K. (1998) *Proc. Natl. Acad. Sci. USA* **95**, 13585–13590. TROSY in triple-resonance experiments: new perspectives for sequential NMR assignment of large proteins.
59. Fernández, C., Hilty, C., Bonjour, S., Adeishvili, K., Pervushin, K. and Wüthrich, K. (2001) *FEBS Lett.* **504**, 173–178. Solution NMR studies of the integral membrane proteins OmpX and OmpA from *Escherichia coli*.
60. Riek, R., Wider, G., Pervushin, K. and Wüthrich, K. (1999) *Proc. Natl. Acad. Sci. USA* **96**, 4918–4923. Polarization transfer by cross-correlated relaxation in solution NMR with very large molecules.

61. Fiaux, J., Bertelsen, E., Horwich, A. and Wüthrich, K. (2002) *Nature* **418**, 207–211. NMR analysis of a 900K GroEL–GroES complex.
62. Pellecchia, M., Sebbel, P., Hermanns, U., Wüthrich, K. and Glockshuber, R. (1999) *Nature Struct. Biol.* **6**, 336–339. Pilus chaperone FimC–adhesin FimH interactions mapped by TROSY-NMR.
63. Frickel, E.M., Riek, R., Jelesarov, I., Helenius, A., Wüthrich, K. and Ellgaard, L. (2002) *Proc. Natl. Acad. Sci. USA* **99**, 1954–1959. TROSY-NMR reveals interaction between ERp57 and the tip of the calreticulin P-domain.
64. Fernández, C., Hilty, C., Wider, G. and Wüthrich, K. (2002) *Proc. Natl. Acad. Sci. USA* **99**, 13533–13537. Lipid–protein interactions in DHPC micelles containing the integral membrane protein OmpX investigated by NMR spectroscopy.